

Vaccine Schedule for South Africa - 2026

The schedule is to be read in conjunction with the footnotes below.

Compiled by Amayeza Info Services - Vaccine Helpline - 0860 160 160

Age of child	EPI schedule	Private
At birth	OPV (0) BCG *See footnotes regarding hep B vaccine at birth	OPV (0) BCG *See footnotes regarding hep B vaccine at birth
6 weeks	OPV (1) Rotarix® (1) PCV10-Cipla® (1) Hexaxim® (1)	OPV (1) Rotarix® (1) or Rotateq® (1) or Rota Vaccine Liquid Cipla®(1) Prevenar 13® or Synflorix® ³ or Vaxneuvance® Hexaxim® (1) or (Infanrix Hexa® (1st dose at 8 weeks)) ¹
10 weeks	Hexaxim® (2)	Rotateq® (2) ² or Rota Vaccine Liquid Cipla®(2) Hexaxim® (2) or (Infanrix Hexa® (2nd dose at 12 weeks)) ¹ Bexsero® (1)
14 weeks	Rotarix® (2) PCV10-Cipla® (2) Hexaxim® (3)	Rotarix® (2) or Rotateq® (3) or Rota Vaccine Liquid Cipla®(3) Prevenar 13® (2) or Synflorix® ³ (2) or Vaxneuvance® Hexaxim® (3) or (Infanrix Hexa® (3rd dose at 16 weeks)) ¹
18 weeks		Bexsero® (2) (At least 8 weeks after first dose of Bexsero®)
6 months	MR SII® ⁴ (1)	Measles vaccine Cipla® ⁴ Influenza Vaccine
9 months	PCV10-Cipla® (3)	Synflorix® or Vaxneuvance® or Prevenar 13® if using 2+1 schedule. (Do not give now if using PCV 3+1 schedule ³) Menactra® (1)
12 months	MR SII® (2) ⁴	Priorix® or Omzyta® (1)
12 - 15 months		Avaxim 80® or Havrix Jnr® (1) Varilrix® ⁶ or Onvara® ⁷ (1) Menactra® (2) (At least 3 months after the first dose of Menactra®) Bexsero® (3)
18 months	Hexaxim® (4)	Hexaxim® (4) or Infanrix Hexa® (4) Avaxim 80® or Havrix Jnr® (2) (At least 6 months after the first dose of Hep A)
6 years	Adacel®	Tetraxim® ⁸ or Boostrix tetra® or Adacel Quadra® Priorix® or Omzyta® (2) Varilrix® (2) ^{6,7} or Onvara® (2)
9 years	Cervarix® ⁹	Cervarix® or Gardasil® or Gardasil 9® ¹⁰ (from 9 years) (Second dose 6 months later)
12 years	Adacel®	Adacel Quadra®, Adacel®, Boostrix® or Boostrix Tetra®

• Hep B SII® (EPI), Engerix B® or Euvax® (0) given at birth to babies born to mothers who test positive for hepatitis B

General:

(0) Birth dose which doesn't count as part of primary series

(1) First dose in a series

(2) Second dose in a series

(3) Third dose in a series

(4) Fourth dose - a booster

Referenced notes (superscripts)

- 1 If the hexavalent vaccine -Infanrix-hexa® is given according to the EPI schedule (6, 10 and 14 weeks), then a birth dose of hepatitis B vaccine is required. If no birth dose is given, then Infanrix Hexa® should only be given from 8 weeks of age. However, if Hexaxim® is used according to the EPI schedule, a birth dose of HBV is NOT required. The first three doses must be at least 4 weeks apart.
- 2 If Rotateq® vaccine or Rota vaccine liquid Cipla® is used, then 3 doses are required, 6, 10 and 14 weeks. If Rotarix® is used, only 2 doses are given – 6 and 14 weeks.
- 3 If PCV is given as a 2+1 schedule, it is given at 6 and 14 weeks and at 9 months. When Synflorix® or PCV10-Cipla® is used as per the 3+1 schedule, it is given at 6, 10 and 14 weeks and the 4th dose should be given at least 6 months after the third dose. When Prevenar 13® Vaxneuvance® (PCV15) is used as per the 3+1 schedule, it is given at 6,10,14 weeks and the 4th dose may be given from 12 months of age (12-15 months).
- 4 MR SII® should not be given with any other vaccines at 6 months of age but can be given with other vaccines from 9 months of age. Cipla measles vaccine may be given on the same day as any other vaccine.
- 5 The first time that a child gets the flu vaccine, 2 doses a month apart are indicated if the child is less than 9 years old.
- 6 Varilrix® can be given any time from 9 months of age but is probably most effective if given over the age of 12 months. If not given on the same day as measles vaccines, must then be separated by at least one month. The second dose can be given as early as 4-6 weeks after the first dose but is routinely given at 5-6 years of age.
- 7 A second dose of Onvara® for children 12 months up to 13 years should be administered at least 3 months following the initial dose. Patients 13 years and older need two doses at least 4 to 8 weeks apart.
- 8 Internationally a DTaP-containing vaccine is recommended at 6 years, but Tdap or Tdap-IPV may also be used.
- 9 Cervarix® in the EPI for girls only. Given to girls >9 years in public schools currently as a single dose vaccination (off label)
- 10 HPV vaccines are for boys and girls. As per PI, for children 9 -13/14 years of age, the course consists of 2 doses six months apart. For older patients give 3 doses as per the PI. Latest information (off label) from WHO states that 1 or 2 doses (6 months apart) are sufficient for patients up to 21 years of age and patients 21 years and older only need two doses given at least 6 months apart. For immunocompromised patients use dosing as per PI.



























Other paediatric vaccines available, for use in certain situations.

Vaxigrip Tetra® / Influvac®, Hiberix® (Hib - Haemophilus influenzae type b), Twinrix® (Hep A paed + hep B adult), Pneumovax 23® (Pneumococcal infection, from 2 years of age for those at high risk) and Stamaril®

References for vaccine schedule:

- a Expanded programme of immunisation (EPI) (From April 2009, 2015, 2022 and 2024)
- b Package inserts: B-OPV, BCG, Infanrix-Hexa, Hexaxim, Prevenar, Varilrix, Gardasil 9, Gardasil, Cervarix, Adacel, Adacel quadra, Synflorix, Boostrix, Boostrix Tetra, Priorix, Rotateq, Rotarix, Influvac, Vaxigrip Tetra, Menactra, Onvara, Omzyta, Measles vaccine Cipla, Hep B SII, MR SII, PCV-10 Cipla, Avaxim, Havrix, Tetraxim, Hiberix, Twinrix, Pneumovax, Vaxneuvance, Bexsero, Rota vaccine Liquid Cipla.
- c Communication with GSK (Upper age limit of Infanrix-Hexa®)
- d CDC pink book <http://www.cdc.gov/vaccines/Pubs/pinkbook/pink-chapters.htm>
- e NAGI (National Advisory Group on Immunisation)

KNOW YOUR PRODUCTS - REGISTERED VACCINES IN SOUTH AFRICA

Vaccine	Abbreviation	Trade name(s) [®]	Age Indications
Diphtheria, tetanus, acellular pertussis and polio vaccine 	DTaP-IPV	Tetraxim [®]	From 6 weeks of age up to 12 years
 Diphtheria, tetanus, acellular pertussis/inactivated polio, <i>Haemophilus influenzae</i> type b and hepatitis B vaccine	DTaP-IPV-Hib-HBV	Hexaxim [®]	From 6 weeks to 5 years
		Infanrix [®] Hexa	From 8 weeks to 3 years
<i>Haemophilus influenzae</i> type b vaccine 	Hib (PRP-T)	Hiberix [®]	Infants from 6 weeks of age
Hepatitis A vaccine 	HepA	Havrix [®] Junior	From 12 months through 15 years
		Avaxim 80 [®]	
		Avaxim 160 [®]	From 16 years through adulthood
Hepatitis B vaccine 	HepB	Engerix-B [®]	From birth through adulthood (age-dependent dose)
		Euvax B [®]	
		HepB SII [®]	
Hepatitis A (paed) and Hepatitis B (adult) combination vaccine 	HepA-HepB	Twinrix [®]	From 12 months through adulthood (age-dependent schedule)
Human papillomavirus vaccine 	HPV	Cervarix [®]	From 9 years and older for both females and males
		Gardasil [®]	Gardasil ages 9 through 45 years (females) Gardasil ages 9 through 26 years (males)
		Gardasil 9 [®]	From 9 years and older for both females and males
Influenza vaccine 	IIV	Influenza vaccine	From 6 months of age (depending on product used)
Measles Vaccine 		Measles Vaccine Cipla [®]	From 9 months of age (EPI from 6 months)
Measles, rubella vaccine 	MR	MR SII [®]	From 9 months of age (EPI from 6 months)
Measles, mumps, and rubella vaccine 	MMR	Priorix [®]	From 12 months to 6 years
		Omzyta [®]	From 12 months of age
Meningococcal serogroups A, C, W, Y vaccine 	MenACWY-D	Menactra [®]	From 9 months of age through 55 years
Meningococcal serogroup B 	4CMenB	Bexsero [®]	From 2 months of age
Pneumococcal conjugate vaccine 	PCV10 (SII)	PCV10-Cipla [®]	From 6 weeks of age up to 2 years (EPI until 6 years)
	PCV10 (GSK)	Synflorix [®]	From 6 weeks to 5 years
	PCV13	Prevenar-13 [®]	From 6 weeks of age through adulthood
	PCV15	Vaxneuvance [®]	From 6 weeks of age through adulthood
Pneumococcal polysaccharide vaccine 	PPSV23	Pneumovax [®] 23	From 2 years of age
Poliovirus vaccine 	OPV	B-OPV [®]	From birth through adulthood (EPI until 6 months)
Rabies vaccine 		Verorab [®]	From birth through adulthood
		Chirorab [®]	
		Rabies vaccine Cipla [®]	
Rotavirus vaccine 	RV1	Rotarix [®]	From 6 weeks until 24 weeks, 0 days
	RV5	RotaTeq [®]	From 6 weeks until 32 weeks, 6 days
	RV5	Rota vaccine Liquid Cipla [®]	From 6 weeks up to one year of age
Respiratory Syncytial Virus vaccine 	RSV	Abrysvo [®]	In pregnancy from week 28 - 36 weeks Patients ≥60 years of age
Tetanus vaccine 	TT	Tetanus Vaccine Cipla [®]	Infants, children, adults
		Tetanus-Toxoid Bio [®]	From 7 years of age
Tetanus and reduced amounts of diphtheria, and acellular pertussis vaccine 	Tdap	Adacel [®]	From 4 years of age
		Boostrix [®]	
Tetanus and reduced amounts of diphtheria, acellular pertussis vaccine and inactivated polio vaccine 	Tdap-IPV	Adacel Quadra [®]	From 3 Years of age
		Boostrix Tetra [®]	From 4 years of age
Tuberculosis vaccine 	BCG	Bacille Calmette Guerin [®]	From birth up to 1 year of age (EPI) TB Guidelines until 10 years
Typhoid vaccine 	Vi-PS	Typhim Vi [®]	From 2 years of age
Varicella vaccine 	VAR	Varilrix [®]	From 9 months of age
		Onvara [®]	From 12 months of age
Yellow Fever vaccine 	YF	Stamaril [®]	From 9 months of age

 Live Vaccine  Inactivated Vaccine